

CHEERLEADING INJURIES



AOSSM SPORTS TIPS

Legend has it that cheerleading started with a University of Minnesota student standing up in the stands and leading his fellow students in “cheering” for their team during a football game. Cheerleading has morphed drastically since that first instance. Today, it’s considered an athletic activity that incorporates elements of dance and gymnastics along with stunts and pyramid formations. In 2002, there were an estimated 3.5 million people in the U.S. participating as cheerleaders, from as young as six-years-old to those who cheerlead for professional athletic teams. While cheerleading is meant to support an athletic team, its intense competitions at the high school and collegiate levels have created a whole new dynamic, including increased risk for injury.

WHAT TYPES OF INJURIES ARE MOST COMMON IN CHEERLEADING?

The Consumer Product Safety Council (CPSC) estimated that 16,000 emergency room visits occurred from cheerleading in 2002 (the latest year for data). While not as frequent as injuries in other sports, cheerleading injuries tend to be more severe. Cheerleading injuries make up more than half of the catastrophic injuries in female athletes. All areas of the body can be affected by cheerleading injuries, but the most commonly affected areas include the wrists, shoulders, ankles, head, and neck.

HOW CAN INJURIES BE PREVENTED?

In an attempt to curb the amount of catastrophic injuries in cheerleading, restrictions have been placed on stunts.

Ranging from height restrictions in human pyramids, to the thrower-flyer ratio, to the number of spotters that must be present for each person lifted above shoulder level. For example: the limit for pyramids is two body lengths at the high school level and 2.5 body lengths for the college level, with the base cheerleader in direct contact with the performing surface. Base supporters must remain stationary and the suspended person is not allowed to be inverted or rotate on dismount.

Basket toss stunts in which a cheerleader is thrown into the air (sometimes as high as 20 feet) are only allowed to have four throwers. The person being tossed (flyer) is not allowed to drop the head below a horizontal plane with the torso. One of the throwers must remain behind the flyer at all times during the toss.

Mats should be used during practice sessions and as much as possible during competitions. Cheerleaders should not attempt a stunt if tired, injured, or ill, as this may disrupt their focus causing the stunt to be performed in an unsafe manner.

The importance of a qualified coach is also critical. Coaching certification is encouraged.

Precautions should always be taken during inclement weather for all stunts. Also, a stunt should not be attempted without proper training, and not until the cheerleader is confident and comfortable with performing the stunt. Supervision should be provided at all times during stunt routines.

As with any sport, proper conditioning and training are important to minimize injury, including:

- Targeting the lower back, stomach, and shoulders to gain strength through resistive exercises.
- Improving flexibility through regular stretching, yoga or pilates instruction

HOW ARE INJURIES TREATED?

One of the more common injuries cheerleaders suffer is an ankle or wrist sprain or strain. For treatment of a sprain or strain, remember RICE:

- R** Rest the injured site for at least 24 hours.
- I** Ice the injury at least every hour for 10–20 minutes during the initial 4 hours after injury. Icing then can be done 10–20 minutes 4 times a day for two days.
- C** Compress the injured site with a snug, elastic bandage for 48 hours.
- E** Elevate the injured limb for at least 24 hours.

Immediate medical attention is required for any cheerleader with a suspected head or neck injury.



Reprinted with permission Boden et al.

References

Campbell, John D, and Barry P. Boden.
Cheerleading Injuries. *Sports Medicine Update*. September/October 2008.

Boden BP, Tacchetti R, Mueller FO.
Catastrophic cheerleading injuries. *Am Journal of Sports Med*. 2003;31 :881-888.

Shields BJ, Smith GA. Cheerleading related injuries to children 5 to 18 years of age: United States, 1990-2002. *Pediatrics*. 2006; 117:122-129.

Sports Tips are brought to you by the American Orthopaedic Society for Sports Medicine. They provide general information only and are not a substitute for your own good judgement or consultation with a physician. To order multiple copies of this fact sheet or learn more about other orthopaedic sports medicine topics, please visit www.sportsmed.org.



The American Orthopaedic
Society for Sports Medicine

Copyright © 2008. American Orthopaedic
Society for Sports Medicine. All rights reserved.
Multiple copy reproduction prohibited.